

Total number of printed pages-8

3 (Sem-5/CBCS) MAT HC 2

2024

**MATHEMATICS**

(Honours Core)

Paper : MAT-HC-5026

**(Linear Algebra)**

Full Marks : 80

Time : Three hours

***The figures in the margin indicate full marks for the questions.***

1. Answer the following questions as directed :  
1×10=10

(a) Give reason why a line in  $\mathbb{R}^2$  not passing through the origin is not a subspace of  $\mathbb{R}^2$ .

(b) Express  $W = \left\{ \begin{bmatrix} 6a-b \\ a+b \\ -7a \end{bmatrix} ; a, b \in \mathbb{R} \right\}$

as span of two vectors.

Contd.



- (c) State whether the following statement is true or false :

"A finite dimensional vector space has exactly one basis."

- (d) Find the dimension of the subspace of all vectors in  $\mathbb{R}^3$  whose first and third entries are equal.
- (e) 0 is an eigenvalue of a matrix  $A$  if and only if  $A$  is \_\_\_\_\_. (Fill in the blank)
- (f) When is a square matrix said to be diagonalizable?
- (g) Write the kernel of the linear transformation  $T : \mathbb{R}^3 \rightarrow \mathbb{R}^3$  such that  $T(x, y, z) = (x, 0, z)$ .
- (h) If  $u = \begin{bmatrix} 2 \\ -5 \\ -1 \end{bmatrix}$  and  $v = \begin{bmatrix} 3 \\ 2 \\ -3 \end{bmatrix}$ , then compute  $u \cdot v$ .
- (i) What is the distance between the vectors  $\vec{u} = (7, 1)$  and  $\vec{v} = (3, 2)$  in the  $\mathbb{R}^2$  plane?

- (j) What do you mean by orthogonal vectors in an inner product space?

2. Answer the following questions:  $2 \times 5 = 10$

- (a) Let  $A = \begin{bmatrix} -6 & 12 \\ -3 & 6 \end{bmatrix}$  and  $w = \begin{bmatrix} 2 \\ 1 \end{bmatrix}$ .

Determine if  $w$  is in null space of  $A$ .

- (b) Let  $\mathbb{P}_3$  be the vector space of all polynomials of degree at most 3.

Are the vectors

$p(t) = 1 + t^2$  and  $q(t) = 1 - t^2$  linearly independent in  $\mathbb{P}_3$ ? Justify your answer.

- (c) Find the eigenvalues of the matrix

$$A = \begin{bmatrix} 4 & 0 & 0 \\ -2 & 1 & 0 \\ 5 & 4 & -3 \end{bmatrix}$$

- (d) Let  $\mathcal{B} = \{b_1, b_2, b_3\}$  be a basis for a vector space  $V$  and  $T : V \rightarrow \mathbb{R}^2$  be a linear transformation such that

$$T(x_1 b_1 + x_2 b_2 + x_3 b_3) = \begin{bmatrix} 2x_1 - 4x_2 + 5x_3 \\ -x_2 + 3x_3 \end{bmatrix}.$$

Find the matrix for  $T$  relative to  $\mathcal{B}$ .



- (e) Let  $v=(1,-2,2,0)$  be a vector in  $\mathbb{R}^4$ . Find a unit vector  $u$  in the same direction as  $v$ .

3. Answer **any four** questions :  $5 \times 4 = 20$

- (a) If a vector space  $V$  has a basis  $\mathcal{B} = \{b_1, b_2, \dots, b_n\}$ , then prove that any set in  $V$  containing more than  $n$  vectors must be linearly dependent.

(b) Let  $b_1 = \begin{bmatrix} 1 \\ -3 \end{bmatrix}$ ,  $b_2 = \begin{bmatrix} 2 \\ -5 \end{bmatrix}$  and  $x = \begin{bmatrix} -2 \\ 1 \end{bmatrix}$ .  
 $3+2=5$

- (i) Show that the set  $\mathcal{B} = \{b_1, b_2\}$  is a basis of  $\mathbb{R}^2$

- (ii) Find the coordinate vector  $[x]$  of  $x$  relative to  $\mathcal{B}$ .

- (c) Given that 2 is an eigenvalue of the

matrix  $A = \begin{bmatrix} 4 & -1 & 6 \\ 2 & 1 & 6 \\ 2 & -1 & 8 \end{bmatrix}$

Find a basis for the corresponding eigenspace.

- (d) Prove that an  $n \times n$  matrix  $A$  is diagonalizable if and only if  $A$  has  $n$  linearly independent eigenvectors.

- (e) Compute the orthogonal projection of  $\begin{bmatrix} 1 \\ 7 \end{bmatrix}$  onto the line through  $\begin{bmatrix} -4 \\ 2 \end{bmatrix}$  and the origin.

- (f) If  $\{u, v\}$  is an orthonormal set in an inner product space  $V$ , then show that  $\|u - v\| = \sqrt{2}$ .

Answer either (a) **or** (b) from each of the following questions :  $10 \times 4 = 40$

4. (a) Find the rank and the nullity of the matrix  $10$

$$A = \begin{bmatrix} -2 & -5 & 8 & 0 & -17 \\ 1 & 3 & -5 & 1 & 5 \\ 3 & 11 & -19 & 7 & 1 \\ 1 & 7 & -13 & 5 & -3 \end{bmatrix}$$

- (b) Let  $b_1 = \begin{bmatrix} -9 \\ 1 \end{bmatrix}$ ,  $b_2 = \begin{bmatrix} -5 \\ -1 \end{bmatrix}$ ,  $c_1 = \begin{bmatrix} 1 \\ -4 \end{bmatrix}$  and

consider the bases for  $\mathbb{R}^2$  given by  $\mathcal{B} = \{b_1, b_2\}$  and  $\mathcal{C} = \{c_1, c_2\}$   $5+5=10$

- (i) Find the change-of-coordinates matrix from  $\mathcal{C}$  to  $\mathcal{B}$



(ii) Find the change-of-coordinates matrix from  $C$  to  $B$

5. (a) (i) If  $n \times n$  matrices  $A$  and  $B$  are similar, then show that they have the same characteristic polynomial. 3

(ii) If  $A = \begin{bmatrix} 4 & 0 & -2 \\ 2 & 5 & 4 \\ 0 & 0 & 5 \end{bmatrix}$  then find an

invertible matrix  $P$  and a diagonal matrix  $D$  such that  $A = PDP^{-1}$  7

(b) (i) Let  $B = \{b_1, b_2, b_3\}$  and  $D = \{d_1, d_2\}$  be bases for vector spaces  $V$  and  $W$  respectively. Let  $T: V \rightarrow W$  be a linear transformation with the property that

$$T(b_1) = 3d_1 - 5d_2, T(b_2) = -d_1 + 6d_2, T(b_3) = 4d_2.$$

Find the matrix for  $T$  relative to  $B$  and  $D$ . 5

(ii) Let  $A$  be a real  $2 \times 2$  matrix with a complex eigenvalue

$$\lambda = a - bi (b \neq 0) \text{ and an associated}$$

eigenvector  $v$  in  $\mathbb{C}^2$ . Show that

$$A(Re v) = a Re v + b Im v \text{ and}$$

$$A(Im v) = b Re v + a Im v \quad 5$$

6. (a) Let,  $B = \left\{ \begin{bmatrix} 2 \\ -5 \\ 1 \end{bmatrix}, \begin{bmatrix} 4 \\ -1 \\ 2 \end{bmatrix} \right\}$  be a basis for a

subspace  $W$  of  $\mathbb{R}^3$ . Using the Gram-Schmidt process construct an orthogonal basis for  $W$ .

Hence, find an orthonormal basis.

$$8+2=10$$



- (b) What do you mean by an inner product on a vector space  $V$ ? Consider the inner product in

$$\mathbb{R}^2 \text{ defined by } \langle u, v \rangle = 4u_1v_1 + 5u_2v_2$$

$$\text{where } u = (u_1, u_2), \quad v = (v_1, v_2) \in \mathbb{R}^2.$$

If  $x = (1, 1)$  and  $y = (5, -1)$ , then find

$\|x\|, \|y\|$  and  $|\langle x, y \rangle|^2$ . Also, show that in an inner product space  $V$  over  $\mathbb{R}$ ,

$$\langle u, v \rangle = \frac{1}{4} \|u + v\|^2 - \frac{1}{4} \|u - v\|^2, \quad \forall u, v \in V.$$

$$2+3+5=10$$

7. (a) State Cayley-Hamilton theorem for matrices. Verify the theorem for the matrix.

$$2+6+2=10$$

$$M = \begin{bmatrix} 2 & -1 & 1 \\ -1 & 2 & -1 \\ 1 & -1 & 2 \end{bmatrix}$$

and hence find  $M^{-1}$ .

- (b) If possible, diagonalize the symmetric matrix

$$10$$

$$A = \begin{bmatrix} 6 & -2 & -1 \\ -2 & 6 & -1 \\ -1 & -1 & 5 \end{bmatrix}$$